AGROMAN 6 L

PREVENTS AND TREATS MANGANESE DEFICIENCY
ALLOWED IN ORGANIC FARMING

AGROMAN 6 L is a specific product whose use is aimed at the prevention and the cure of plant physiological disorders related to manganese deficiency. Manganese is a catalyser of oxidative processes in the plant. It is involved in respiration, photosynthesis and it has a role in controlling auxin metabolism together with other microelements. It also favors buds fertility, fruit-set and it increases resistance to cold.

CROP	TIME OF APPLICATION	DOSE/HECTARE*
Grapes	At pre-flowering and pea-sized berries, 1-2 applications every 8-10 days	2-3 Kg
Kiwifruit, Stone fruits (Plum, Peach, Nectarine, Cherry, Apricot) e Pome fruits (Pear, Apple, Quince)	At early vegetative phases, at pre-flowering or at the appearance of first deficiency symptoms, 2-3 applications every 8-10 days	2-3 Kg
Olive	At vegetative restart, 2 applications every 8-10 days	2-3 Kg
Citrus (Tangerine, Lemon, Clementine, Bergamot, Orange)	At pre-flowering or at the appearance of first deficiency symptoms	2-3 Kg
Fruiting vegetables (Pumpkin, Zucchini, Tomato, Pepper, Melon, Eggplant, Cucumber, Watermelon)	From early vegetative phases or at the appearance of first deficiency symptoms	2-3 Kg
Cereal crops (Triticale, Sorghum, Rye, Rice, Barley, Corn, Wheat, Oats)	From early vegetative phases or at the appearance of first deficiency symptoms	2-3 Kg
Beets	At vegetative restart and when leaf canopy meets between the rows, 2 applications every 8-10 days	2-3 Kg
Industrial crops (Tobacco, Soybeans, Industrial tomato, Sunflower, Cotton, Rapeseed, Sugarcane)	From early vegetative phases or at the appearance of first deficiency symptoms	2-3 Kg
Flowers and ornamentals	At post-emergence or post-transplanting, to be repeated at pre-flowering	2-3 Kg

COMPOSITION	
Manganese (Mn) soluble in water	6%
Manganese (Mn) chelated by EDTA	6%

PHYSICO-CHEMICAL FEATURES	
LIQUID	
pH (sol 1%)	6.0
Conductivity E.C. µS/cm (1‰)	192
Density (g/cm^3)/Specific weight	1.26

PACKAGING: 1 - 6 - 12 KG

The choice of the dose is subordinated to various factors and can be varied when necessary. All applications can be repeated in relation to the different crop needs. You can contact our Technical Service for the correct application on specific soils and under specific climate conditions.